

Chemistry 271, Section 22xx

Your Name: \_\_\_\_\_

Prof. Jason Kahn

University of Maryland, College Park

Your SID #: \_\_\_\_\_

General Chemistry and Energetics

Your Section #: \_\_\_\_\_

Exam I (100 points total)

October 12, 2011

You have 50 minutes for this exam.

Exams written in pencil or erasable ink will not be re-graded under any circumstances.

Explanations should be concise and clear. I have given you more space than you should need. There is extra space on the last page if you need it.

You will need a calculator for this exam. No other study aids or materials are permitted.

Partial credit will be given, *i.e.*, if you don't know, guess.

Useful Equations:

$$K_a = [\text{H}^+][\text{A}^-]/[\text{HA}]$$

$$\text{pH} = -\log([\text{H}^+])$$

$$K_b = [\text{BH}^+][\text{HO}^-]/[\text{B}]$$

$$F = ma$$

$$e^{i\pi} + 1 = 0$$

$$PV = nRT$$

$$K_w = [\text{H}^+][\text{HO}^-] = 10^{-14}$$

$$\text{pH} = \text{p}K_a + \log([\text{A}^-]/[\text{HA}])$$

$$\text{pH (e.p.)} = (\text{p}K_{a1} + \text{p}K_{a2})/2$$

$$R = 0.08206 \text{ L}\cdot\text{atm}/\text{mole K}$$

$$0 \text{ }^\circ\text{C} = 273.15 \text{ K}$$

$$\text{p}K_a = -\log(K_a)$$

$$K_p = K_c(\text{RT})^{\Delta n}$$

$$P^2/a^3 = 4\pi^2/MG$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Honor Pledge: At the end of the examination time, please write out the following sentence and sign it, or talk to me about it:

“I pledge on my honor that I have not given or received any unauthorized assistance on this examination.”

+1 point extra credit for filling in this box

**1. (20 pts) Short Answer (2 pts each)**pH of 1 M HNO<sub>3</sub>\_\_\_\_\_

pOH of 0.1 M NaOH\_\_\_\_\_

pOH of 0.1 M HOAc (pK<sub>a</sub> 4.75) + 0.1 M NaOAc\_\_\_\_\_

pH of 0.001 M NaOH\_\_\_\_\_

pK<sub>a</sub> + pK<sub>b</sub> = \_\_\_\_\_ (for K<sub>a</sub> and K<sub>b</sub> referring to a conjugate acid/base pair)[H<sup>+</sup>] at pH 7\_\_\_\_\_[OH<sup>-</sup>] at pH 9\_\_\_\_\_When  $Q < K$ , the reaction will proceed (circle one)      forward.      backward.

The Henderson-Hasselbach relationship is always \_\_\_\_\_, sometimes \_\_\_\_\_.

The pH at the first equivalence point of a polyprotic acid titration is given by \_\_\_\_\_.

Score for the page\_\_\_\_\_

**2. (30 pts) Acid-Base Equilibria**

Consider the pH obtained upon dissolving a weak monoprotic acid HA in water, as a function of its total concentration  $C_0$  and its  $K_a$ . This is a problem you have done many times, here we are exploring a general formula. The equilibrium is of course



(a; 12 pts) Initially, assume that “ $x$ ” can be ignored in the denominator and show that  $\text{pH} = \frac{1}{2} \log(K_a \times C_0)$

(b; 8 pts) Now, repeat the problem but do not assume that “ $x$ ” is small, i.e. use the quadratic formula to derive a general formula for the pH


(c; 4 pts) Show that the more complicated expression you just obtained reduces to the simpler expression from (a) if  $C_0 \gg K_a$ . (This gives us a more precise description of exactly when  $x$  is negligible in the denominator.)

- (d; 6 pts) Physically, what happens to the % Dissociation of a weak acid as  $K_a$  increases or  $C_0$  decreases?  
Give an explanation for the  $C_0$  effect based on LeChatelier's principle or the dynamic balance of rates.

### 3. (20 pts) Buffers and Titration

The graph shows seven titrations at progressively lower concentrations of a weak acid titrated with strong base. The  $x$  axis is in terms of equivalents of base added relative to the acid, so the actual concentration of base added is also decreasing as  $C_0$  decreases.

As usual, we ignore dilution during each individual titration, so the total acid  $[HA] + [A^-]$  is constant.



- (a; 6 pts) What is the  $pK_a$  of the weak acid being used, and how do you know?

(b; 4 pts) Why does the pH at the equivalence point decrease as  $C_0$  decreases? A qualitative answer is fine.

(c; 4 pts) The graph illustrates the critical features of buffers. How does it show us the utility of “10X” or “100X” reaction buffers in the lab?

(d; 6 pts) Explain why the pH at the end (1.2 equivalents, top right corner) of the  $C_0 = 0.5$  M titration is at pH 13.

**4. (10 pts) pH effects on Enzymes**

The proposed mechanism shown is the essence of catalysis by aspartyl proteases, a class of enzymes that includes HIV protease. They have classic bell-shaped pH rate profiles like those we have seen in class, with pKa's typically at around 3 and 5.




Fig. 3 Proposed catalytic mechanism for aspartyl proteases.

(a; 10 pts) Which residue is associated with the pKa of 3, and what is its function in the mechanism? In other words, why does the reaction fail when it is performed at a pH much below 3? Which residue has pKa = 5, and what is its function? Why does the reaction fail at basic pH?

**5. (20 pts) Equilibrium Manipulations.**

We will confirm that any acid on the left column of the table can protonate any base that is below it in the right column. For example, formic acid,  $\text{HCOOH}$ ,  $\text{pK}_a = 3.75$ , should be able to protonate sulfite,  $\text{SO}_3^{2-}$ . The  $\text{pK}_a$  of hydrogen sulfite ( $\text{HSO}_3^-$ ) is 6.97.


(a; 3 pts) What is the net reaction for formic acid protonating sulfite?

(b; 7 pts) Write down the base dissociation reaction for sulfite and calculate its  $\text{pK}_b$  and  $K_b$ .

(c; 3 pts) Add the acid dissociation reaction of formic acid and the base dissociation reaction of sulfite.

(d; 3 pts) What other equilibrium do we need to add to give us the net reaction from part a?

(e; 4 pts) Calculate the overall equilibrium reaction constant for the reaction of part a.



Chemistry: The Science in Context 3/e Figure 17.4  
© 2012 W. W. Norton & Company, Inc.

Page	Score
1	/1
2	/20
3	/15
4	/13
5	/12
6	/18
7	/12
<b>Total</b>	<b>/101</b>

Score for the page \_\_\_\_\_